

Kleinian groups of small Hausdorff dimension are classical Schottky groups. I

Yong Hou

dedicated to Ying Zhou

Abstract

It has been conjectured that the Hausdorff dimensions of nonclassical Schottky groups are strictly bounded from below. In this first part of our works on this conjecture, we prove that there exists a universal positive number $\lambda > 0$, such that any 2-generated Kleinian groups with limit set of Hausdorff dimension $< \lambda$ are classical Schottky groups.

1 Introduction and Main Theorem

Let $\mathbb{H}^3$ be the hyperbolic 3-space. A subgroup Γ of $\mathrm{PSL}(2, \mathbb{C}) = \mathrm{Iso}(\mathbb{H}^3)$ is called a Kleinian group if it is discrete. Let $x \in \mathbb{H}^3$. The orbit of x under action of Γ is denoted by Γx . The limit set Λ_Γ of Γ is defined as $\Lambda_\Gamma = \overline{\Gamma x} \cap \partial \mathbb{H}^3$. By definition, Λ_Γ is smallest closed Γ -invariant subset of $\partial \mathbb{H}^3$. The group Γ is called second kind if $\Lambda_\Gamma \neq \partial \mathbb{H}^3$, otherwise it is said to be first kind. The set $\Omega_\Gamma = \partial \mathbb{H}^3 - \Lambda_\Gamma$ is the region of discontinuity, and Γ acts properly discontinuously on Ω_Γ .

Let $\{\Delta_1, \Delta'_1, \dots, \Delta_k, \Delta'_k\}$ be a collection of disjoint closed Jordan curves in the Riemann sphere $\overline{\mathbb{C}}$ and D_i, D'_i the topological disk bounded by Δ_i and Δ'_i respectively. Suppose we have $\{\gamma_i\}_1^k \subset \mathrm{PSL}(2, \mathbb{C})$ such that $\gamma_i(\Delta_i) = \Delta'_i$ and $\gamma_i(D_i) \cap D'_i = \emptyset$, then the group Γ generated by $\{\gamma_1, \dots, \gamma_k\}$ is a free Kleinian group of rank- k , and Γ is called marked Schottky group with markings $\{\gamma_1, \dots, \gamma_k\}$. A finitely generated Kleinian group Γ is called Schottky group if its marked Schottky group for some markings. If there exists a generating set $\{\gamma_i, \dots, \gamma_k\}$ such that all Δ_i, Δ'_i can be taken as circles then it is

called a marked classical Schottky group with classical markings $\{\gamma_i, \dots, \gamma_k\}$, and $\{\gamma_1, \dots, \gamma_k\}$ are called classical generators. A Schottky group Γ is called classical Schottky group if there exists a classical markings for Γ . For a Schottky group Γ , the manifold $\mathbb{H}^3/\Gamma$ is homeomorphic to the interior of a handle body of genus k . We denote by $\mathfrak{J}_k$ the set of all rank- k Schottky groups, and $\mathfrak{J}_{k,o}$ the set of all rank- k *classical* Schottky groups. One simply way to topologies $\mathfrak{J}_k$ into a topological space is by identify it with the space of moduli for Riemann surface of genus k . It is known that not all Schottky groups are classical Schottky groups, in-fact the space of classical Schottky groups is not even dense in the space of Schottky groups [14], also see [9].

In [9], Peter Doyle proved that there exists a universal upper bound for all finitely generated classical Schottky groups. It was originally Phillips and Sarnak in [16] proved that there exists a universal upper bound for all classical Schottky groups of dimension greater than 3. Hence, our result can be considered as a converse to these results.

Let $\mathfrak{D}_\Gamma$ denote the Hausdorff dimension of Λ_Γ . Our main result is,

Theorem 1.1. *There exists a universal $\lambda > 0$, such that any 2-generated Kleinian group Γ with $\mathfrak{D}_\Gamma < \lambda$ is a classical Schottky group.*

Note that our result can be viewed as the converse of the result by Doyle [9] and Phillips and Sarnak [16].

We prove Theorem 1.1 by using result of [12], and an selection process of generators. The proof is divided into three main steps.

To lead up to the proof, we first do some preliminary estimates on the locations of the fixed points of a generating set of a given Schottky group. These estimates gives us a sufficient control of how the fixed-points-set of a set of generators change in terms of the Hausdorff dimension of the limit set of the group. The main ingredient of the proofs of these estimates relies on the result of [12], Theorem 1.1 rewritten in the trace form.

Next we obtain an set of sufficient conditions for any given sequence of Schottky groups to contain a subsequence of classical Schottky groups in the unit ball hyperbolic space. These conditions are stated in the upper-half space hyperbolic model. The idea is that in the unit ball model of hyperbolic space if the radius of isometric circles of the generators decreases sufficiently faster than the reduction of the gaps between any of the fixed points of the generators then it will eventually becomes classical generators. This is done by first transform the generators, with one of the generator (in the proof we take this generator to be the one of shortest translation length) put into

vertically passing through the origin with fixed points north and south poles, and then these generators are projected into upper-half space.

In the first step of the proof, we consider Schottky subspaces of the Schottky space that are consists of Schottky groups Γ for which there exists a generating set S_Γ for Γ with the set of fixed points of S_Γ on the boundary sphere on the unit 3-ball hyperbolic space are mutually bounded away by a given constant $Z > 0$. In this step we prove that Theorem 1.1 holds for these Schottky subspaces. This is proved via contradiction. Suppose Γ_n is a sequence of nonclassical Schottky groups in the subspace with Hausdorff dimension of Λ_{Γ_n} decreasing to 0. The idea is that we first transform these generators of the generating sets S_{Γ_n} into the standard form with the generator of S_{Γ_n} of shortest translation length put in vertically. If no generator of S_{Γ_n} is of bounded translation length when $\mathfrak{D}_{\Gamma_n} \rightarrow 0$ then its easy to see it will lead to a contradiction. On the other hand, its a simply corollary of [12] that there can exists at most one generator of bounded translation length per S_{Γ_n} when $\mathfrak{D}_{\Gamma_n} \rightarrow 0$. If such a generator do exists, then we first make a careful change of generators which will be constructed based on estimates that its fixed-points set will be “minimally excluded” from isometric spheres of the generator of bounded translation length. Working with these generating sets we will show that with appropriate additional transformations and changes the generators with translation length that is not bounded will always grow sufficiently fast to lead to disjoint isometric spheres and is also disjoint from rest of isometric spheres of other generators. One of the crucial tool in estimating this growth is the strong form of the inequality of [12]. As we will show that when one of the generator is of bounded translation length then it will force the needed growth for the rest of the generators. And with appropriate choice of generators this will lead to a classical Schottky groups.

The second step, we prove that on the nonclassical Schottky space there exists a universal lower bounds for $\mathfrak{D}_\Gamma + Z_\Gamma$. Essentially means that we can not have simultaneous arbitrary small of Hausdorff dimension and minimal gaps of its fixed-points set on the space of nonclassical Schottky groups. This is proved using generator selection and result of step one.

In the last step, we prove that when $\mathfrak{D}_\Gamma$ is taken sufficiently small then Γ can be taken as Schottky group. This is based on basic topological arguments and some well known results on Kleinian groups. Although we proved (easy standard topological argument) that all sufficiently small Hausdorff dimension of the limit set of an finitely generated Kleinian group is a Schottky group, but as Dick Canary pointed out that its simpler in the 2-generated

case based on the result of Peter Shalen which assures that any 2-generated Kleinian group is either free or confinite volume.

The paper is organized as follows: Section 2, we state a strong form of Theorem 1.1 given in [12] for two generators, which we shall use for selecting generators. Section 3, for a given sequence of Schottky groups Γ_n with bounds on Z_Γ (see section 3), we prove inequalities that will enable us to control fixed points of a given sequence of generators of Γ_n in relation to the Hausdorff dimensions $\mathfrak{D}_n$ of Γ_n . These will be used in the selection processes. Section 4, a sufficient condition for a given pair of generator to be classical generator is proved which we will use in our generator selection process. Section 5, we will use this in our generator selection process and prove that Schottky groups with small $\mathfrak{D}_n$ and bounds on Z_Γ are classical Schottky groups. Section 6, we prove result that will remove the bound condition on Z_Γ . Section 7 completes the proofs of our main theorem.

2 Free Group Actions

Given a Kleinian group Γ , the critical exponent of Γ is the unique positive number D_Γ such that the Poincaré series of Γ given by $\sum_{\gamma \in \Gamma} e^{-s \text{dist}(x, \gamma x)}$ is divergent if $s < D_\Gamma$ and convergent if $s > D_\Gamma$. If the Poincaré series diverges at $s = D_\Gamma$ then Γ is said to be divergent. Bishop-Jones showed that $D_\Gamma \leq \mathfrak{D}_\Gamma$ for all analytically finite Kleinian groups Γ . In-fact, if Γ is topologically tame ($\mathbb{H}^3/\Gamma$ homeomorphic to the interior of a compact manifold-with-boundary) then $D_\Gamma = \mathfrak{D}_\Gamma$. The critical exponent is a geometrically rigid object in the sense that decrease in D_Γ corresponds to decrease in geometric complexity.

Next we state the following theorem in [12], which provides the relation between the group action and the critical exponent.

Theorem 2.1 (Hou[12]). *Let Γ be a free Kleinian group of rank- k with free generating set $\mathcal{S}$, and $x \in \mathbb{H}^3$ then*

$$\sum_{\gamma \in \mathcal{S}} \frac{1}{1 + \exp(D_\Gamma \text{dist}(x, \gamma x))} \leq \frac{1}{2}.$$

In particular we have at least $k-1$ distinct elements $\{\gamma_{i_j}\}_{1 \leq j \leq k-1}$ of $\mathcal{S}$ that satisfies $\text{dist}(x, \gamma_{i_j} x) \geq \log(3)/D_\Gamma$, and at least one element γ_{i_j} with $\text{dist}(x, \gamma_{i_j} x) \geq \log(2k-1)/D_\Gamma$.

The following is a useful corollary of Theorem 2.1 stated here for the case of Γ is a free group of rank-2.

Corollary 2.2. *Let $\mathcal{S} = \{\gamma_1, \gamma_2\}$ be a generating set for a free group Γ . Let $x \in \mathbb{H}^3$. Then*

$$\text{dist}(x, \gamma_2 x) \geq \frac{1}{D_\Gamma} \log \left(\frac{e^{D_\Gamma \text{dist}(x, \gamma_1 x)} + 3}{e^{D_\Gamma \text{dist}(x, \gamma_1 x)} - 1} \right)$$

Corollary 2.3. *Let $\mathcal{S} = \{\gamma_1, \gamma_2\}$ be a generating set for a free group Γ . Let $x \in \mathbb{H}^3$. Assume $\text{dist}(x, \gamma_1 x) \geq \text{dist}(x, \gamma_2 x)$. Let m be any integer. Then at least one of the element γ' of $\mathcal{S}' = \{\gamma_1^m \gamma_2, \gamma_1^{m+1} \gamma_2\}$ satisfies $\text{dist}(x, \gamma' x) \geq \log 3 / D_\Gamma$.*

For a loxodromic element $\gamma \in \text{PSL}(2, \mathbb{C})$ we denote the axis of γ in $\mathbb{H}^3$ by α_γ . Define $\beta_\mathcal{S}$ the axial-distance function of a finitely generated purely loxodromic $\Gamma \subset \text{PSL}(2, \mathbb{C})$ with generating set $\mathcal{S}$ as $\beta_\mathcal{S} := \max_{\gamma_i, \gamma_j \in \mathcal{S}} \text{dist}(\alpha_{\gamma_i}, \alpha_{\gamma_j})$. Next proposition although is not used subsequently, but is of interest in its own right. The proof is trivial.

Proposition 2.4. *Let $x \in \mathbb{H}^3$. Let $R > 0$ be any positive real numbetr. Let Γ be a finitely generated free purely loxodromic Kleinian group. Then there is a free generating set $\mathcal{S}$ such that $\min_{\gamma \in \mathcal{S}} \text{dist}(x, \gamma x) \geq \log 3 / D_\Gamma$ and $\beta_\mathcal{S} < R$.*

Corollary 2.5. *Let $\mathcal{S} = \{\gamma_1, \dots, \gamma_k\}$ be a generating set for a free group Γ . Let $x \in \mathbb{H}^3$. Assume there is $\gamma_i \in \mathcal{S}, i \neq 1$ such that $\text{dist}(x, \gamma_1 x) \geq \text{dist}(x, \gamma_i x)$. Then there exists some $\{a_i\}_{i=2}^k$ nonnegative integers with $\sum_i a_i \leq 1$ such that every elements γ' of the generating set $\mathcal{S}' = \{\gamma_1, \gamma_1^{a_2} \gamma_2, \dots, \gamma_1^{a_k} \gamma_k\}$ satisfies $\text{dist}(x, \gamma' x) \geq \log 3 / D_\Gamma$.*

3 Trace, Fixed Points and Hausdorff Dimension

In this section we study the relation of fixed points of a given free generating set of a free Kleinian group with Hausdorff dimension of its limit set. What we like to do is to find a relationship of the distribution of fixed points of one generator in terms of the growth of the other generators and Hausdorff dimension of its limit set. By having this type of relations we will be able to construct new set of generators from the given generating set with prescribed

distribution of its fixed points. The new set of generators will be a crucial ingredience in the proof of our theorem.

We will assume the following throughout this section. Set $D_n = D_{\Gamma_n}$. Let $\{\Gamma_n\}$ be a sequence of purely loxodromic free group with $D_n \rightarrow 0$ generated by $\alpha_n, \beta_n \in \text{PSL}(2, \mathbb{C})$ in the upper space model $\mathbb{H}^3$ with $\alpha_n = \begin{pmatrix} \lambda_n & 0 \\ 0 & \lambda_n^{-1} \end{pmatrix}$, $|\lambda_n| > 1$. Set $\beta_n = \begin{pmatrix} a_n & b_n \\ c_n & d_n \end{pmatrix}$. Set the notation, $\eta_n = -\frac{d_n}{c_n}, \zeta_n = \frac{a_n}{c_n}$. In general, for a given Mobius transformation g with fixed points $\neq$ to ∞ . we set the following notations throught rest of the paper.

Notation 1. we set the following notations throught rest of the paper.

- $\mathfrak{R}_g$ the radius of its isometric circles.
- $\eta_g = -\frac{d_g}{c_g}, \zeta_g = \frac{a_g}{c_g}$.
- $\{z_{g,l}, z_{g,u}\}$ and $\{z_{g,-}, z_{g,+}\}$ are the two fixed points of g in $\mathbb{C}$, with $|z_{g,l}| \leq |z_{g,u}|$ and $z_{g,\pm}$ given by quadratic formula with subscripts $\pm$ corresponds to $\pm\sqrt{\text{tr}^2(g) - 4}$.
- $\mathcal{L}_g$ is the axis of g .
- T_g is the translation length of g .

Set $Z_{\beta_n} := \min\{|z_{\beta_n,-} - z_{\beta_n,+}|, |\frac{1}{z_{\beta_n,-}} - \frac{1}{z_{\beta_n,+}}|\}$, and

$$Z_{<\alpha_n, \beta_n>} := \min\{Z_{\beta_n}, |z_{\beta_n,+}|, |z_{\beta_n,-}|, |z_{\beta_n,+}|^{-1}, |z_{\beta_n,-}|^{-1}\}.$$

For $\epsilon > 0$, we say that $Z_{\Gamma} > \epsilon$, if there exists a generating set $<\alpha, \beta>$ of Γ such that $Z_{<\alpha, \beta>} > \epsilon$.

We assume throughout this section that there exists $M, \Delta > 0$ such that $Z_{<\alpha_n, \beta_n>} \geq \Delta > 0$ and $T_{\alpha_n} < M$ for all n .

In the following, denote by k_n, l_n to be any integers that satisfies the constraint: $T_{\alpha_n^{k_n}}, T_{\alpha_n^{l_n}} < M$.

Given any two sequences of real numbers $\{p_n, q_n\}$, we use the notation $p_n \asymp q_n$ iff, there exists $\sigma > 0$ such that, $\sigma^{-1} < \liminf \frac{p_n}{q_n} \leq \limsup \frac{p_n}{q_n} < \sigma$.

First we will state Corollary 2.2 in the trace form. The proof is stright forward and is left as a exercises for the reader.

Proposition 3.1. *There exists $\rho > 0$ depends on Δ , such that*

$$|\text{tr}(\beta_n)| \geq \rho \left(\frac{|\lambda_n|^{2D_n} + 3}{|\lambda_n|^{2D_n} - 1} \right)^{\frac{1}{2D_n}}$$

for large n .

Notation 2. Given sequences $\{p_n\}, \{q_n\}$ of numbers we write:

$$|z_{g_n, \pm} - p_n| < q_n$$

if there exists N such that for each $n > N$ we have one of the following holds,

(i)

$$|z_{g_n, +} - p_n| < q_n$$

(ii)

$$|z_{g_n, -} - p_n| < q_n$$

Lemma 3.2. *There exists constant $\rho > 0$ such that,*

$$\left| z_{\alpha_n^{k_n} \beta_n \alpha_n^{l_n}, \pm} - \zeta_{\alpha_n^{k_n} \beta_n \alpha_n^{l_n}} \right| + \left| z_{\alpha_n^{k_n} \beta_n \alpha_n^{l_n}, \mp} - \eta_{\alpha_n^{k_n} \beta_n \alpha_n^{l_n}} \right| < \frac{\rho}{|\operatorname{tr}(\beta_n)| |\lambda_n^{l_n - k_n}|}$$

for large n .

Remark 3.3. Lemma 3.2 is a estimates of how fast the fixed points converges, its not important to our applications in this paper that which fixed point converges to η_g and which converges to ζ_g for a given $g \in \operatorname{PSL}(2, \mathbb{C})$. However a more precise statement of Lemma 3.2 which dichotomizes the above inequality for large n would be:

$$(i) \quad \sqrt{\operatorname{tr}^2(\alpha_n^{k_n} \beta_n \alpha_n^{l_n})} = \operatorname{tr}(\alpha_n^{k_n} \beta_n \alpha_n^{l_n})$$

$$\left| z_{\alpha_n^{k_n} \beta_n \alpha_n^{l_n}, +} - \zeta_{\alpha_n^{k_n} \beta_n \alpha_n^{l_n}} \right| + \left| z_{\alpha_n^{k_n} \beta_n \alpha_n^{l_n}, -} - \eta_{\alpha_n^{k_n} \beta_n \alpha_n^{l_n}} \right| < \frac{\rho}{|\operatorname{tr}(\beta_n)| |\lambda_n^{l_n - k_n}|}$$

$$(ii) \quad -\sqrt{\operatorname{tr}^2(\alpha_n^{k_n} \beta_n \alpha_n^{l_n})} = \operatorname{tr}(\alpha_n^{k_n} \beta_n \alpha_n^{l_n})$$

$$\left| z_{\alpha_n^{k_n} \beta_n \alpha_n^{l_n}, -} - \zeta_{\alpha_n^{k_n} \beta_n \alpha_n^{l_n}} \right| + \left| z_{\alpha_n^{k_n} \beta_n \alpha_n^{l_n}, +} - \eta_{\alpha_n^{k_n} \beta_n \alpha_n^{l_n}} \right| < \frac{\rho}{|\operatorname{tr}(\beta_n)| |\lambda_n^{l_n - k_n}|}$$

Corollary 3.4. *For any $\delta > 0$ there exists $\epsilon > 0$ such that if $D_n < \epsilon$ then,*

$$\left| z_{\alpha_n^{k_n} \beta_n \alpha_n^{l_n}, \pm} - \zeta_{\alpha_n^{k_n} \beta_n \alpha_n^{l_n}} \right| + \left| z_{\alpha_n^{k_n} \beta_n \alpha_n^{l_n}, \mp} - \eta_{\alpha_n^{k_n} \beta_n \alpha_n^{l_n}} \right| < \delta(|\lambda_n|^2 - 1).$$

The same conditions for dichotomy decomposition of the inequality as given in Remark 3.3.

Proof. Let us assume that $\sqrt{\text{tr}^2(\alpha_n^{k_n} \beta_n \alpha_n^{l_n})} = \text{tr}(\alpha_n^{k_n} \beta_n \alpha_n^{l_n})$. Since $Z_{\langle \alpha_n, \beta_n \rangle} \geq \Delta$, there exists positive numbers $0 < c_1(\Delta) < c_2(\Delta), c_3(\Delta) < c_4(\Delta), c_5(\Delta) < c_6(\Delta)$ such that for large n , $c_1 < |\frac{\sqrt{\text{tr}^2(\beta_n) - 4}}{2c_n}| < c_2$, $c_3 < |(a_n + b_n)/2c_n| < c_4$ and $c_5 < |\frac{a_n}{c_n}|, |\frac{d_n}{c_n}| < c_6$. In particular, for $D_n \rightarrow 0$ Theorem 2.1 implies $|\text{tr}(\beta_n)| \rightarrow \infty$, and $|\text{tr}(\beta_n)| \asymp |c_n|$ for large n . Therefore the fixed points $z_{+,-}(\beta_n)$ of β_n must $\rightarrow \{\frac{a_n}{c_n}, -\frac{d_n}{c_n}\}$. There exists $c, c', N > 0$ such that:

$$\begin{aligned} \left| z_{\alpha_n^{k_n} \beta_n \alpha_n^{l_n}, +} - \frac{a_n}{c_n} \lambda_n^{2k_n} \right| &= \left| \frac{(a_n \lambda_n^{k_n + l_n} - d_n \lambda_n^{-l_n - k_n}) + \sqrt{\text{tr}^2(\alpha_n^{k_n} \beta_n \alpha_n^{l_n}) - 4}}{2c_n \lambda_n^{l_n - k_n}} - \frac{a_n \lambda_n^{k_n + l_n}}{c_n \lambda_n^{l_n - k_n}} \right| \\ &= \left| \frac{\sqrt{\text{tr}^2(\alpha_n^{k_n} \beta_n \alpha_n^{l_n}) - 4} - \sqrt{\text{tr}^2(\alpha_n^{k_n} \beta_n \alpha_n^{l_n})}}{2c_n \lambda_n^{l_n - k_n}} \right| \\ &\leq c \frac{1}{|\text{tr}(\beta_n)|} \left| \sqrt{\text{tr}^2(\alpha_n^{k_n} \beta_n \alpha_n^{l_n}) - 4} - \sqrt{\text{tr}^2(\alpha_n^{k_n} \beta_n \alpha_n^{l_n})} \right| \\ &\leq c \frac{|\sqrt{\text{tr}^2(\alpha_n^{k_n} \beta_n \alpha_n^{l_n}) - 4}|}{|\text{tr}(\beta_n)|} \left| \frac{1}{\sqrt{1 - 4/\text{tr}^2(\alpha_n^{k_n} \beta_n \alpha_n^{l_n})}} - 1 \right| \\ &\leq \frac{c'}{|\text{tr}(\beta_n)|} \quad \text{for } n > N. \end{aligned}$$

Since for sufficient large n there exists $a, b > 0$, $e^{T_{\beta_n}} \leq a|\text{tr}(\beta_n)|^2$ and $e^{T_{\alpha_n^{k_n} \beta_n \alpha_n^{l_n}}} \leq b|\text{tr}(\alpha_n^{k_n} \beta_n \alpha_n^{l_n})|^2$, we therefore need $\min\{c'ae^{-T_{\beta_n}}, c'be^{-T_{\alpha_n^{k_n} \beta_n \alpha_n^{l_n}}}\} < \delta(|\lambda_n|^2 - 1)$. By $|\lambda_n|^2 = e^{T_{\alpha_n}}$, we require T_{β_n} to satisfy

$$\max\{T_{\beta_n}, T_{\alpha_n^{k_n} \beta_n \alpha_n^{l_n}}\} \geq 2 \log \left(\frac{\frac{c'}{\delta}}{e^{T_{\alpha_n}} - 1} \right). \quad (1)$$

Hence it follows there exists $\rho > 0$ such that if $D_n < \rho$ then, T_{β_n} satisfies the above inequality. Therefore we have $|z_{\beta_n, +} - \frac{a_n}{c_n}| < \delta(|\lambda_n|^2 - 1)$ for $D_n < \rho$. The proof for the other part is same.

The case $-\sqrt{\text{tr}^2(\alpha_n^{k_n} \beta_n \alpha_n^{l_n})} = \text{tr}(\alpha_n^{k_n} \beta_n \alpha_n^{l_n})$ is similar with $z_{\alpha_n^{k_n} \beta_n \alpha_n^{l_n}, +}$ replace with $z_{\alpha_n^{k_n} \beta_n \alpha_n^{l_n}, -}$ and vice versa. $\square$

Corollary 3.5. *There exists constants $\delta, \rho > 0$ such that, if $|\text{tr}(\alpha_n^{k_n} \beta_n \alpha_n^{l_n})| \rightarrow \infty$ then,*

$$\left| z_{\alpha_n^{k_n} \beta_n \alpha_n^{l_n}, \pm} - \zeta_{\alpha_n^{k_n} \beta_n \alpha_n^{l_n}} \right| + \left| z_{\alpha_n^{k_n} \beta_n \alpha_n^{l_n}, \mp} - \eta_{\alpha_n^{k_n} \beta_n \alpha_n^{l_n}} \right| < \frac{\rho}{|\text{tr}(\alpha_n^{k_n} \beta_n \alpha_n^{l_n})| |\text{tr}(\beta_n)|}$$

for all $D_n < \delta$. The same conditions for dichotomy decomposition of the inequality as given in Remark 3.3.

Proof. Let us assume that $\sqrt{\text{tr}^2(\alpha_n^{k_n} \beta_n \alpha_n^{l_n})} = \text{tr}(\alpha_n^{k_n} \beta_n \alpha_n^{l_n})$. It follows from the proof of Corollary 3.4,

$$\begin{aligned} \left| z_{\alpha_n^{k_n} \beta_n \alpha_n^{l_n}, +} - \frac{a_n}{c_n} \lambda_n^{2k_n} \right| &\leq c \frac{|\sqrt{\text{tr}^2(\alpha_n^{k_n} \beta_n \alpha_n^{l_n})} - 4|}{|\text{tr}(\beta_n)|} \left| \frac{1}{\sqrt{1 - 4/\text{tr}^2(\alpha_n^{k_n} \beta_n \alpha_n^{l_n})}} - 1 \right| \\ &< \frac{\rho}{|\text{tr}(\alpha_n^{k_n} \beta_n \alpha_n^{l_n})| |\text{tr}(\beta_n)|} \end{aligned}$$

The proof for the other part is same.

The case $-\sqrt{\text{tr}^2(\alpha_n^{k_n} \beta_n \alpha_n^{l_n})} = \text{tr}(\alpha_n^{k_n} \beta_n \alpha_n^{l_n})$ is similar with $z_{\alpha_n^{k_n} \beta_n \alpha_n^{l_n}, +}$ replace with $z_{\alpha_n^{k_n} \beta_n \alpha_n^{l_n}, -}$ and vice versa. $\square$

Lemma 3.6. *Let α_n, β_n as in Lemma 3.4. Let k_n, l_n be integers such that $T_{\alpha_n}, T_{\alpha_n^{k_n}}, T_{\alpha_n^{l_n}} < M$ for all n . Then there exists a constant $\sigma_1, \sigma_2 > 0$ such that,*

$$\frac{\sigma_1 |\text{tr}(\alpha_n^{k_n} \beta_n \alpha_n^{l_n})|}{|\text{tr}(\beta_n)|} \geq |z_{\alpha_n^{k_n} \beta_n \alpha_n^{l_n}, +} - z_{\alpha_n^{k_n} \beta_n \alpha_n^{l_n}, -}| \geq \frac{\sigma_2 |\text{tr}(\alpha_n^{k_n} \beta_n \alpha_n^{l_n})|}{|\text{tr}(\beta_n)|},$$

for all n sufficiently large.

Proof. Note that,

$$\begin{aligned} \left| \frac{a_n}{c_n} \lambda_n^{2k_n} - \left(-\frac{d_n}{c_n} \lambda_n^{-2l_n} \right) \right| &= \left| \frac{a_n \lambda_n^{k_n+l_n} + d_n \lambda_n^{-k_n-l_n}}{c_n \lambda_n^{l_n-k_n}} \right| = \left| \frac{\text{tr}(\alpha_n^{k_n} \beta_n \alpha_n^{l_n})}{c_n \lambda_n^{l_n-k_n}} \right| \quad (2) \\ &\geq \left| \frac{\text{tr}(\alpha_n^{k_n} \beta_n \alpha_n^{l_n})}{\rho \sqrt{\text{tr}^2(\beta_n) - 4}} \right| \quad (3) \end{aligned}$$

The last inequality follows from $c_1 \leq \sqrt{\text{tr}(\beta_n) - 4}/2c_n \leq c_2$ (using the notation as in the proof of Lemma 3.4) and $T_{\alpha_n^{k_n}} < M$. Now the result follows from Corollary 3.4. $\square$

Lemma 3.7. *Let α_n, β_n as in Lemma 3.4. Let k_n, l_n be integers such that $T_{\alpha_n}, T_{\alpha_n^{k_n}}, T_{\alpha_n^{l_n}} < M$ for all n . Then for at least one $i \in \{0, 1\}$, and any integers k'_n, l'_n with $|(k_n - k'_n) + (l_n - l'_n)| = i$, we have $|z_{\alpha_n^{k'_n} \beta_n \alpha_n^{l'_n}, +} - z_{\alpha_n^{k'_n} \beta_n \alpha_n^{l'_n}, -}| \geq \frac{\kappa}{D_n |\text{tr}(\beta_n)|}$, for all n sufficiently large.*

4 Sufficient Conditions

Let $(\mathbb{B}, \text{dist}_B)$ be the unit ball model of the hyperbolic 3-space. Let $\pi : \mathbb{B} \longrightarrow \mathbb{H}^3$ be the stereographic hyperbolic isometry.

Given a loxodromic element, α of $\text{PSL}(2, \mathbb{C})$ acting on the unit ball $\mathbb{B}$ hyperbolic 3-space, denote by $S_{\alpha, r}$ and $S_{\alpha^{-1}, r}$ the isometric spheres of euclidean radius r of α . We set $\lambda_{\pi_*(\alpha)}$ as the multiplier of $\pi_*(\alpha)$ in the upper space model $\mathbb{H}^3$. For $R > 0$, set C_R as the circle in $\mathbb{C}$ about origin of radius R .

Proposition 4.1. *Let α be a loxodromic element of $\text{PSL}(2, \mathbb{C})$ acting on $\mathbb{B}$, with axis passing through the origin and fixed points on north and south poles. Then $\pi(S_{\alpha, r} \cap \partial\mathbb{B}), \pi(S_{\alpha^{-1}, r} \cap \partial\mathbb{B})$ maps to $C_{\frac{1}{\lambda_{\pi_*(\alpha)}}}, C_{\lambda_{\pi_*(\alpha)}}$.*

Proof. Let T_α be the translation length of α . The euclidean radius r is given by $r^{-1} = \sinh(\frac{T_\alpha}{2})$. In terms of $\lambda_{\pi_*(\alpha)}$,

$$r = \frac{2|\lambda_{\pi_*(\alpha)}|}{|\lambda_{\pi_*(\alpha)}|^2 - 1}$$

Set $\mathbf{e} = (0, 0, 1)$ as the north pole of $\partial\mathbb{B}$. Let $\delta_\alpha, \delta_{\alpha^{-1}} > 0$ denote the radius of $\pi(S_{\alpha, r} \cap \partial\mathbb{B})$ and $\pi(S_{\alpha^{-1}, r} \cap \partial\mathbb{B})$ respectively. Then for $x \in S_{\alpha, r} \cap \partial\mathbb{B}$, δ_α is given by,

$$\begin{aligned} \delta_\alpha^2 &= \frac{4r^2}{(1+r^2)|x - \mathbf{e}|^4} \\ \text{and} \quad |x - \mathbf{e}|^2 &= \frac{4}{|\lambda_{\pi_*(\alpha)}|^2 + 1} \\ \text{this implies} \quad \delta_\alpha &= |\lambda_{\pi_*(\alpha)}|. \end{aligned}$$

Similarly for $x \in S_{\alpha^{-1}, r} \cap \partial\mathbb{B}$ we have,

$$\begin{aligned} \delta_{\alpha^{-1}}^2 &= \frac{4r^2}{(1+r^2)|x - \mathbf{e}|^4} \\ \text{and} \quad |x - \mathbf{e}|^2 &= \frac{4|\lambda_{\pi_*(\alpha)}|^2}{|\lambda_{\pi_*(\alpha)}|^2 + 1} \\ \text{which gives} \quad \delta_{\alpha^{-1}} &= \frac{1}{|\lambda_{\pi_*(\alpha)}|}. \end{aligned}$$

□

Lemma 4.2. *Let $\langle \alpha_n, \beta_n \rangle$ be generators for Schottky groups Γ_n in upper space model $\mathbb{H}^3$ with $|\operatorname{tr}(\beta_n)| \rightarrow \infty$ and $\alpha_n = \begin{pmatrix} \lambda_n & 0 \\ 0 & \lambda_n^{-1} \end{pmatrix}$, $|\lambda_n| > 1$. Suppose either there exists $\Lambda > 1$, such that for large n , we have $|\lambda_n| < \Lambda$ and,*

- $|\lambda_n|^{-1} < |z_{\beta_n, l}| \leq |z_{\beta_n, u}| < |\lambda_n|$, and
-

$$\liminf_n \left\{ \frac{1}{(|z_{\beta_n, u}| - |\lambda_n|) |\operatorname{tr}(\beta_n)|}, \frac{1}{(|z_{\beta_n, l}| - |\lambda_n|^{-1}) |\operatorname{tr}(\beta_n)|} \right\} = 0,$$

or there exists $\kappa > 0$ and for large n , we have $|\lambda_n| > \kappa$ and,

- $\kappa^{-1} < |z_{\beta_n, l}| \leq |z_{\beta_n, u}| < \kappa$, and
-

$$\liminf_n \left\{ \frac{1}{(|z_{\beta_n, u}| - \kappa) |\operatorname{tr}(\beta_n)|}, \frac{1}{(|z_{\beta_n, l}| - \kappa^{-1}) |\operatorname{tr}(\beta_n)|} \right\} = 0,$$

then there exists a subsequence such that for i large, $\pi^{-1} \langle \alpha_{n_i}, \beta_{n_i} \rangle \pi$ are classical generators for Γ_{n_i} in the unit ball model $\mathbb{B}$.

Proof. Let us suppose there exists a subsequence $\langle \alpha_{n_i}, \beta_{n_i} \rangle$ that satisfies the first set of conditions. First assume that for large i , $|z_{\beta_{n_i}, u} - z_{\beta_{n_i}, l}| > \delta > 0$.

Let r_i, ρ_i denote the euclidean radius of the isometric spheres of $\pi^{-1} \alpha_{n_i} \pi$ and $\pi^{-1} \beta_{n_i} \pi$ respectively. There exists $\delta_1 > 0$ such that $\rho_i \leq \delta_1 |\operatorname{tr}(\beta_{n_i})|^{-1}$. This implies for $x_i \in C_{|z_{\beta_{n_i}, u}|}$ and $y_i \in C_{|\lambda_{n_i}|}$,

$$\frac{\rho_i}{|\pi^{-1} x_i - \pi^{-1} y_i|} \leq \frac{\sqrt{2} \delta_1}{|\operatorname{tr}(\beta_{n_i})| |\pi^{-1} x_i - \mathbf{e}| |\pi^{-1} y_i - \mathbf{e}| |x_i - y_i|}.$$

Since $|\lambda_{n_i}| < \Lambda$, there exists $\delta_2 > 0$, such that

$$\lim_i \frac{\rho_i}{|\pi^{-1} x_i - \pi^{-1} y_i|} \leq \lim_i \frac{\sqrt{2} \delta_1 \delta_2}{|\operatorname{tr}(\beta_{n_i})| |x_i - y_i|} = 0.$$

Similarly there exists $\delta_3 > 0$, such that for $w_i \in C_{|z_{\beta_{n_i}, l}|}$ and $z_i \in C_{|\lambda_{n_i}|^{-1}}$,

$$\lim_i \frac{\rho_i}{|\pi^{-1} w_i - \pi^{-1} z_i|} \leq \lim_i \frac{\sqrt{2} \delta_1 \delta_3}{|\operatorname{tr}(\beta_{n_i})| |w_i - z_i|} = 0.$$

Hence it follows that for large i , and Proposition 4.1, the isometric spheres $S_{\pi^{-1}\alpha_{n_i}\pi, r_i}, S_{\pi^{-1}\alpha_{n_i}^{-1}\pi, r_i}, S_{\pi^{-1}\beta_{n_i}\pi, \rho_i}, S_{\pi^{-1}\beta_{n_i}^{-1}\pi, \rho_i}$ are disjoint.

Next let us assume that $|z_{\beta_{n_i}, u} - z_{\beta_{n_i}, l}| \rightarrow 0$. Then for i large, there exists $\delta_4 > 0$ such that $\rho_i \leq \delta_4 |\pi^{-1}z_{\beta_{n_i}, u} - \pi^{-1}z_{\beta_{n_i}, l}| |\text{tr}(\beta_{n_i})|^{-1}$. Hence we have $\rho_i / |\pi^{-1}z_{\beta_{n_i}, u} - \pi^{-1}z_{\beta_{n_i}, l}| \rightarrow 0$, and similar to above estimates and Proposition 4.1, implies $S_{\pi^{-1}\alpha_{n_i}\pi, r_i}, S_{\pi^{-1}\alpha_{n_i}^{-1}\pi, r_i}, S_{\pi^{-1}\beta_{n_i}\pi, \rho_i}, S_{\pi^{-1}\beta_{n_i}^{-1}\pi, \rho_i}$ are disjoint.

The second part of the proposition can be proved similarly. $\square$

5 Schottky Subspaces $\mathfrak{J}_k(\tau)$

For $\tau > 0$, define $\mathfrak{J}_k(\tau) := \{\Gamma \in \mathfrak{J}_k | Z_\Gamma > \tau\}$. Recall that $\mathfrak{J}_k$ denotes set of all Schottky groups of rank k .

Theorem 5.1. *Let $\mathfrak{J}_2$ be the set of all 2-generated Schottky groups. For each $\tau > 0$ there exists a $\nu > 0$ such that $\{\Gamma \in \mathfrak{J}_2(\tau) | D_\Gamma \leq \nu\} \subset \mathfrak{J}_{k,o}$.*

Proof. We prove by contradiction. Assume there exists a sequence $\{\Gamma_n\} \subset \mathfrak{J}_2(\tau)$ of nonclassical Schottky groups with $D_n \rightarrow 0$. By pass to subsequence, we may assume $D_n \rightarrow 0$ monotonically. Set $\Gamma_n = \langle \alpha_n, \beta_n \rangle$ with $Z_{\langle \alpha_n, \beta_n \rangle} > \tau$. We arrange the generators so that $|\text{tr}(\alpha_n)| \leq |\text{tr}(\beta_n)|$. There are two possibilities: (I) There exists a subsequence such that $|\text{tr}(\alpha_{n_i})| \rightarrow \infty$, (II) $|\text{tr}(\alpha_n)| < M$, for some $M > 0$.

Case (I) is trivial. Since both $|\text{tr}(\alpha_{n_i})|, |\text{tr}(\beta_{n_i})| \rightarrow \infty$ as $n \rightarrow \infty$, it follows from $Z_{\langle \alpha_n, \beta_n \rangle} > \tau$, there must exists N such that $\langle \alpha_n, \beta_n \rangle$ becomes classical Schottky groups for $n > N$. A contradiction.

Now we consider case (II). We work in upper space model $\mathbb{H}^3$. Conjugate $\langle \alpha_n, \beta_n \rangle$ by Mobius transformation into $\alpha_n = \begin{pmatrix} \lambda_n & 0 \\ 0 & \lambda_n^{-1} \end{pmatrix}$ with $|\lambda_n| > 1$. Denote $\beta_n = \begin{pmatrix} a_n & b_n \\ c_n & d_n \end{pmatrix}$. It follows from Proposition 3.1, $|\text{tr}(\beta_n)| \rightarrow \infty$. In addition, by conjugation with Mobius transformations, we can assume β_n have $\eta_{\beta_n} = 1$. By replace β_n with β_n^{-1} if necessary, we can assume $|\zeta_n| \leq |\eta_n|$.

Since $Z_{\langle \alpha_n, \beta_n \rangle} > \tau$, there exists $\Delta_1, \Delta_2, \Delta_3, \Delta_4 > 0$ such that, $\Delta_1 < |z_{\beta_n, l}| \leq |z_{\beta_n, u}| < \Delta_2$, and $\Delta_3 < |z_{\beta_n, l} - z_{\beta_n, u}| < \Delta_4$. It follows from Lemma 3.2, $\Delta_1 < |\zeta_n| \leq |\eta_n| < \Delta_2$, and $\Delta_2 < \lim_n |\zeta_n - \eta_n| < \Delta_3$.

For each n , choose integers k_n such that:

$$1 \leq |\zeta_n \lambda_n^{2k_n}| < |\lambda_n^2|.$$

We consider the generating set $\langle \alpha_n, \alpha_n^{k_n} \beta_n \rangle$.

By pass to subsequence if necessary, there are three cases need to be considered (To simplify the notation, we denote subsequences by the same index notation):

- (A) $|\zeta_n \lambda_n^{2k_n}| \rightarrow 1$,
- (B) $||\zeta_n \lambda_n^{2k_n}| - |\lambda_n|^2| \rightarrow 0$,
- (C) case (A), (B) do not occur, and $|\zeta_n \lambda_n^{2k_n}|$, converges in the annulus $\{z | 1 < |z| < |\lambda_n|^2\}$.

5.1 Case (C)

We use the same notation index for subsequences. Since cases (A), (B) do not occur, for large n there exists $1 < |\lambda| < |\lambda_n|$, $\sigma < 1$, such that $|\zeta_n \lambda_n^{2k_n}| \rightarrow \sigma |\lambda|^2$. Let ψ_n be Mobius transformation that fixes $\{0, \infty\}$ and definite by $\psi(x) = \frac{x}{\sqrt{\sigma}|\lambda|}$, $x \in \mathbb{C}$. Then $|\eta_{\psi \alpha_n^{k_n} \beta_n \psi^{-1}}| = \frac{1}{\sqrt{\sigma}|\lambda|}$ and $|\zeta_{\psi \alpha_n^{k_n} \beta_n \psi^{-1}}| \rightarrow \sqrt{\sigma}|\lambda|$. It follows from Lemma 3.2, for n large,

$$\max\{|z_{\psi \alpha_n^{k_n} \beta_n \psi^{-1}, \mp} - \eta_{\psi \alpha_n^{k_n} \beta_n \psi^{-1}}|, |z_{\psi \alpha_n^{k_n} \beta_n \psi^{-1}, \pm} - \zeta_{\psi \alpha_n^{k_n} \beta_n \psi^{-1}}|\} < \frac{\rho}{|\text{tr}(\beta_n)|}$$

Hence by applying Proposition 3.1 to the generators $\langle \psi \alpha_n \psi^{-1}, \psi \alpha_n^{k_n} \beta_n \psi^{-1} \rangle$, implies $|\text{tr}(\psi \alpha_n^{k_n} \beta_n \psi^{-1})| \rightarrow \infty$. We can easily verify that $\langle \psi \alpha_n \psi^{-1}, \psi \alpha_n^{k_n} \beta_n \psi^{-1} \rangle$ satisfies second set of conditions of Lemma 4.2, and so by Lemma 4.2, these will be classical generators for large n , a contradiction.

5.2 case (A)

By pass to a subsequence if necessary, we have two possibilites:

- (A₁) $|\lambda_n|^2 - 1$ is monotonically decreasing to 0.
- (A₂) There exists $\lambda > 1$, such that $|\lambda_n| \geq |\lambda|$ for large n .

5.2.1 (A₁)

Here we either have

$$(i) \quad \limsup_n \left| |\zeta_{\alpha_n^{k_n} \beta_n}| - |\lambda_n|^2 \right| \left| \zeta_{\alpha_n^{k_n} \beta_n} - 1 \right|^{-1} < \infty$$

or

$$(ii) \quad \liminf_n \left| |\zeta_{\alpha_n^{k_n} \beta_n}| - |\lambda_n|^2 \right| \left| \zeta_{\alpha_n^{k_n} \beta_n} - 1 \right|^{-1} \rightarrow \infty.$$

If $\alpha_n^{k_n} \beta_n$ satisfies (ii), we conjugate $\alpha_n^{k_n} \beta_n$ with Mobius transformation ψ_n definite by, $\psi_n(x) = \frac{x}{\zeta_{\alpha_n^{k_n} \beta_n \alpha_n^{l_n}}}$. Consider $(\psi_n \alpha_n^{k_n} \beta_n \alpha_n^{l_n+1} \psi_n^{-1})^{-1}$. Then

$$\limsup_n \left| |\zeta_{\psi_n^{-1} \alpha_n^{-l_n-1} \beta_n^{-1} \alpha_n^{-k_n} \psi_n}| - |\lambda_n|^2 \right| \left| \zeta_{\psi_n \alpha_n^{-l_n-1} \beta_n^{-1} \alpha_n^{-k_n} \psi_n} - 1 \right|^{-1} < \infty.$$

The generator $(\psi_n \alpha_n^{k_n} \beta_n \alpha_n^{l_n+1} \psi_n^{-1})^{-1}$ satisfies (i). Hence replacing the generators if necessary we can always assume the generators satisfies (i). And with out lost of generality we will assume that $\langle \alpha_n, \alpha_n^{k_n} \beta_n \rangle$ satisfies (i).

Consider (i).

In this case, we have either:

$$(i_1) \quad \limsup_n \left| |\zeta_{\alpha_n^{k_n} \beta_n}| - |\lambda_n|^2 \right| \left| \zeta_{\alpha_n^{k_n} \beta_n} - 1 \right|^{-1} > \delta > 0,$$

or

$$(i_2) \quad \limsup_n \left| |\zeta_{\alpha_n^{k_n} \beta_n}| - |\lambda_n|^2 \right| \left| \zeta_{\alpha_n^{k_n} \beta_n} - 1 \right|^{-1} = 0.$$

5.2.1.1 Consider (i₁).

Lemma 5.2. *We have $|z_{\alpha_n^{k_n} \beta_n, -} - z_{\alpha_n^{k_n} \beta_n, +}| \asymp (|\lambda_n|^2 - 1)$, and there exists $c > 0$ such that,*

$$\text{dist}(\mathcal{L}_{\alpha_n}, \mathcal{L}_{\alpha_n^{k_n} \beta_n}) < c \log\left(\frac{1}{|\lambda_n|^2 - 1}\right).$$

Proof. It follows from Lemma 3.2 and our assumption, we only need to show that $\frac{1}{|\text{tr}(\beta_n)|(|\lambda_n|^2 - 1)} \rightarrow 0$. This can be seen from Proposition 3.1,

$$\begin{aligned} \lim \frac{1}{|\text{tr}(\beta_n)|(|\lambda_n|^2 - 1)} &\leq \lim \rho \left(\frac{|\lambda_n|^{2D_n} - 1}{(|\lambda_n|^{2D_n} + 3)(|\lambda_n|^2 - 1)^{2D_n}} \right)^{\frac{1}{2D_n}} \\ &\leq \lim \rho'(|\lambda_n|^2 - 1)^{\frac{1-2D_n}{D_n}} = 0. \end{aligned}$$

□

Lemma 5.3.

$$\lim_n \frac{1}{|\operatorname{tr}(\alpha_n^{k_n} \beta_n)|(|\lambda_n|^2 - 1)} = 0$$

Proof. It follows from Proposition 3.1 and Lemma 5.2, there exists $\rho > 0$ such that,

$$|\operatorname{tr}(\alpha_n^{k_n} \beta_n)| \geq \rho \left(\frac{(|\lambda_n|^{2D_n} + 3)(|\lambda_n|^2 - 1)^{2D_n}}{|\lambda_n|^{2D_n} - 1} \right)^{\frac{1}{2D_n}}.$$

hence we have,

$$\begin{aligned} \lim \frac{1}{|\operatorname{tr}(\alpha_n^{k_n} \beta_n)|(|\lambda_n|^2 - 1)} &\leq \lim \rho \left(\frac{|\lambda_n|^{2D_n} - 1}{(|\lambda_n|^{2D_n} + 3)(|\lambda_n|^2 - 1)^{4D_n}} \right)^{\frac{1}{2D_n}} \\ &\leq \lim \rho' (|\lambda_n|^2 - 1)^{\frac{1-4D_n}{2D_n}} = 0. \end{aligned}$$

□

Hence there exists Mobius transformations $\psi_n(x) = \epsilon_n(x)$ such that the generators, $\langle \psi_n \alpha_n \psi_n^{-1}, \psi_n \alpha_n^{k_n} \beta_n \psi_n^{-1} \rangle$ satisfies conditions of Lemma 4.2.

5.2.1.2 Consider (i_2) .

There exists $|\rho_{\alpha_n^{k_n-1} \beta_n}| \rightarrow 1$ such that $\zeta_{\alpha_n^{k_n-1} \beta_n} \rho_{\alpha_n^{k_n-1} \beta_n} = \eta_{\alpha_n^{k_n-1} \beta_n} = 1$. If $\limsup_n |\rho_{\alpha_n^{k_n-1} \beta_n} - 1| > 0$, then (with the same index notation for subsequence there exists a subsequence) such that

$$\liminf_n |\zeta_{\alpha_n^{k_n-1} \beta_n} - \eta_{\alpha_n^{k_n-1} \beta_n}| > 0.$$

This implies by Lemma 3.2, $\liminf_n |z_{\alpha_n^{k_n-1} \beta_n, +} - z_{\alpha_n^{k_n-1} \beta_n, -}| > 0$. Hence by Proposition 3.1, there exists $\rho > 0$, such that for large n ,

$$|\operatorname{tr}(\alpha_n^{k_n-1} \beta_n)| \geq \rho \left(\frac{|\lambda_n|^{2D_n} + 3}{|\lambda_n|^{2D_n} - 1} \right)^{\frac{1}{2D_n}}.$$

In particular, we have $|\operatorname{tr}(\alpha_n^{k_n-1} \beta_n)| \rightarrow \infty$. Since

$$\lim_n \frac{|\rho_{\alpha_n^{k_n-1} \beta_n}| - 1}{|\lambda_n|^2 - 1} = 0,$$

and $|\operatorname{tr}(\alpha_n^{k_n-1} \beta_n)(|\lambda_n|^2 - 1)| \rightarrow \infty$, it follows that for large n ,

- $|\lambda_n|^{-1} < |\zeta_{\alpha_n^{k_n-1}\beta_n}| \leq |\eta_{\alpha_n^{k_n-1}\beta_n}| < |\lambda_n|$, and
- $\lim_n \left\{ \frac{1}{|\operatorname{tr}(\alpha_n^{k_n-1}\beta_n)|(|\eta_{\alpha_n^{k_n-1}\beta_n}| - |\lambda_n|)}, \frac{1}{|\operatorname{tr}(\alpha_n^{k_n-1}\beta_n)|(|\zeta_{\alpha_n^{k_n-1}\beta_n}| - |\lambda_n|^{-1})} \right\} = 0$.

Hence by Lemma 3.2, $\langle \alpha_n, \alpha_n^{k_n-1}\beta_n \rangle$ satisfies Lemma 4.2.

Consider the case that, $\rho_{\alpha_n^{k_n-1}\beta_n} \rightarrow 1$.

Let ψ_n be the Mobius transformations given by,

$$\psi_n(x) = \frac{1}{\Delta_n} \frac{z_{\alpha_n^{k_n-1}\beta_n,-}x - z_{\alpha_n^{k_n-1}\beta_n,-}z_{\alpha_n^{k_n-1}\beta_n,+}}{x - z_{\alpha_n^{k_n-1}\beta_n,-}}, \quad x \in \mathbb{C}.$$

where $\Delta_n = \frac{1}{z_{\alpha_n^{k_n-1}\beta_n,-}z_{\alpha_n^{k_n-1}\beta_n,+} - z_{\alpha_n^{k_n-1}\beta_n,-}^2}$. Let ϕ_n be the Mobius transformations defined by, $\phi_n(x) = \eta_{\psi_n\alpha_n^{k_n}\beta_n\psi_n^{-1}}^{-1}x$. Then after some computation $\zeta_{\phi_n\psi_n\alpha_n^{k_n}\beta_n\psi_n^{-1}\phi_n^{-1}}$ is given by,

$$\zeta_{\phi_n\psi_n\alpha_n^{k_n}\beta_n\psi_n^{-1}\phi_n^{-1}} = \left(\frac{1 - z_{\alpha_n^{k_n-1}\beta_n,+}}{1 - z_{\alpha_n^{k_n-1}\beta_n,-}} \right) \left(\frac{\alpha_n^{k_n}\beta_n(z_{\alpha_n^{k_n-1}\beta_n,-} - z_{\alpha_n^{k_n-1}\beta_n,+})}{\alpha_n^{k_n}\beta_n(z_{\alpha_n^{k_n-1}\beta_n,+} - z_{\alpha_n^{k_n-1}\beta_n,-})} \right)$$

Lemma 5.4. *We claim that $|\zeta_{\phi_n\psi_n\alpha_n^{k_n}\beta_n\psi_n^{-1}\phi_n^{-1}}| \rightarrow 1$. More precisely,*

$$\left| \left| \zeta_{\phi_n\psi_n\alpha_n^{k_n}\beta_n\psi_n^{-1}\phi_n^{-1}} \right| - 1 \right| \asymp \frac{1}{|\operatorname{tr}(\beta_n)(|\lambda_n|^2 - 1)|}.$$

Proof. It follows from Lemma 3.2 and our assumption, the first factor of the above equation for $\zeta_{\phi_n\psi_n\alpha_n^{k_n}\beta_n\psi_n^{-1}\phi_n^{-1}}$ certainly holds for the lemma. Hence its sufficient to show that the second factor also satisfies the bounds.

After computation the upper bound is given by,

$$\begin{aligned} \lim \left| \frac{\alpha_n^{k_n}\beta_n(z_{\alpha_n^{k_n-1}\beta_n,-}) - z_{\alpha_n^{k_n-1}\beta_n,+}}{\alpha_n^{k_n}\beta_n(z_{\alpha_n^{k_n-1}\beta_n,+}) - z_{\alpha_n^{k_n-1}\beta_n,-}} \right| &\leq \lim \frac{|\zeta_{\alpha_n^{k_n}\beta_n} - z_{\alpha_n^{k_n-1}\beta_n,+}| + |\operatorname{tr}(\beta_n)|^{-1}}{|\zeta_{\alpha_n^{k_n}\beta_n} - z_{\alpha_n^{k_n-1}\beta_n,-}| - |\operatorname{tr}(\beta_n)|^{-1}} \\ &\leq \lim \frac{\left| \frac{|\zeta_{\alpha_n^{k_n}\beta_n} - z_{\alpha_n^{k_n-1}\beta_n,+}|}{|\lambda_n|^2 - 1} + |\operatorname{tr}(\beta_n)(|\lambda_n|^2 - 1)|^{-1} \right|}{\left| \frac{|\zeta_{\alpha_n^{k_n}\beta_n} - z_{\alpha_n^{k_n-1}\beta_n,-}|}{|\lambda_n|^2 - 1} - |\operatorname{tr}(\beta_n)(|\lambda_n|^2 - 1)|^{-1} \right|} \\ &\leq 1 + \delta \lim \frac{1}{|\operatorname{tr}(\beta_n)(|\lambda_n|^2 - 1)|}, \quad \delta > 0. \end{aligned}$$

Similarly we also have lower bound given by $1 - \delta' |\operatorname{tr}(\beta_n)(|\lambda_n|^2 - 1)|^{-1}$ for large n . Hence the lemma follows. $\square$

In the following we set, $\tilde{\alpha}_n = \phi_n \psi_n \alpha_n^{k_n-1} \beta_n \psi_n^{-1} \phi_n^{-1}$ and $\tilde{\beta}_n = \phi_n \psi_n \alpha_n^{k_n} \beta_n \psi_n^{-1} \phi_n^{-1}$. Let $\tilde{\lambda}_n$, denote the multipliers of $\tilde{\alpha}_n$. There are two cases we need to consider on $\tilde{\lambda}_n$, either $\limsup |\tilde{\lambda}_n| > 1$, or $|\tilde{\lambda}_n| \rightarrow 1$.

5.2.1.2a $\limsup |\tilde{\lambda}_n| > 1$.

Let $\langle \tilde{\alpha}_{n_i}, \tilde{\beta}_{n_i} \rangle$ be a subsequence such that $\lim_i |\tilde{\lambda}_{n_i}| > 1$. By Lemma 5.4, $|\zeta_{\tilde{\beta}_{n_i}} - \eta_{\tilde{\beta}_{n_i}}| \rightarrow 0$ and $\eta_{\tilde{\beta}_{n_i}} = 1$, it follows from Lemma 3.2, $z_{\tilde{\beta}_{n_i},+}, z_{\tilde{\beta}_{n_i},-} \rightarrow 1$. Since $|\operatorname{tr}(\tilde{\beta}_{n_i})| \rightarrow \infty$, so its easy to check that $\langle \tilde{\alpha}_{n_i}, \tilde{\beta}_{n_i} \rangle$ satisfies conditions of Lemma 4.2, hence by this Lemma, we have classical generators in $\mathbb{B}$ for large i .

5.2.1.2b $|\tilde{\lambda}_n| \rightarrow 1$, and we proceed with estimate of lower bounds of $|\tilde{\lambda}_n|$.

Lemma 5.5. *Let $\tilde{\lambda}_n$ be the multiplier of $\alpha_n^{k_n-1} \beta_n$ with $|\tilde{\lambda}_n| \rightarrow 1$. Then there exists $\mathcal{N}, \sigma > 0$ such that, $||\tilde{\lambda}_n|^2 - 1| > \sigma |\operatorname{tr}(\beta_n)(|\lambda_n|^2 - 1)|^{-4D_n}$, for $n > \mathcal{N}$.*

Proof. Use matrix representation we can write, $\tilde{\beta}_n = \begin{pmatrix} \tilde{a}_n & \tilde{b}_n \\ \tilde{d}_n & -\tilde{d}_n \end{pmatrix}$. It follows from Lemma 5.4 and $|\operatorname{tr}(\tilde{\beta}_n)| \asymp |\operatorname{tr}(\beta_n)(|\lambda_n|^2 - 1)|$ that,

$$|z_{\tilde{\beta}_n,+} - z_{\tilde{\beta}_n,-}| \asymp |\zeta_{\tilde{\beta}_n} - \eta_{\tilde{\beta}_n}| \asymp |\operatorname{tr}(\beta_n)(|\lambda_n|^2 - 1)|^{-1}.$$

Since $|z_{\tilde{\beta}_n,+} - z_{\tilde{\beta}_n,-}| \asymp |\operatorname{tr}(\beta_n)(|\lambda_n|^2 - 1)| |\tilde{d}_n|^{-1}$, we have $|\tilde{d}_n| \asymp |\operatorname{tr}(\beta_n)(|\lambda_n|^2 - 1)|^2$.

For large n , let $m_n > 0$ be integers defined as,

$$1 + \frac{1}{m_n + 1} \leq |\tilde{\lambda}_n| \leq 1 + \frac{1}{m_n}.$$

From this definition we have, $\lim_n |\tilde{\lambda}_n|^{m_n} = \lim_n (1 + m_n^{-1})^{m_n} = e$. Then by

Lemma 5.4, there exists $N, \delta > 0$ such that for $n > N$ we have,

$$\begin{aligned} |\zeta_{\tilde{\alpha}_n^{m_n} \tilde{\beta}_n} - \eta_{\tilde{\alpha}_n^{m_n} \tilde{\beta}_n}| &\leq \left| |\zeta_{\tilde{\beta}_n}| |\tilde{\lambda}_n|^{m_n} + 1 \right| \\ &\leq \delta(e + 1) \end{aligned}$$

and,

$$\begin{aligned} |\zeta_{\tilde{\alpha}_n^{m_n} \tilde{\beta}_n} - \eta_{\tilde{\alpha}_n^{m_n} \tilde{\beta}_n}| &\geq \left| |\zeta_{\tilde{\beta}_n}| |\tilde{\lambda}_n|^{m_n} - 1 \right| \\ &\geq \delta(e - 1). \end{aligned}$$

Hence it follows from Lemma 3.2, there exists $\kappa > 0$ such that,

$$\kappa^{-1} < |z_{\tilde{\alpha}_n^{m_n} \tilde{\beta}_n, +} - z_{\tilde{\alpha}_n^{m_n} \tilde{\beta}_n, -}| < \kappa.$$

Since,

$$\begin{aligned} |z_{\tilde{\alpha}_n^{m_n} \tilde{\beta}_n, +} - z_{\tilde{\alpha}_n^{m_n} \tilde{\beta}_n, -}| &= \frac{\left| \sqrt{\text{tr}^2(\tilde{\alpha}_n^{m_n} \tilde{\beta}_n) - 4} \right|}{|\tilde{d}_n \tilde{\lambda}_n^{m_n}|} \\ &\asymp \frac{|\text{tr}(\tilde{\alpha}_n^{m_n} \tilde{\beta}_n)|}{|\text{tr}(\tilde{\beta}_n)|^2}, \end{aligned}$$

this implies that $|\text{tr}(\tilde{\alpha}_n^{m_n} \tilde{\beta}_n)| \asymp |\text{tr}(\tilde{\beta}_n)|^2$.

By Corollary 2.2 and above bounds on $|z_{\tilde{\alpha}_n^{m_n} \tilde{\beta}_n, +} - z_{\tilde{\alpha}_n^{m_n} \tilde{\beta}_n, -}|$, there exists $R > 0, \varsigma_n > 0$ with $R^{-1} < \varsigma_n < R$, such that for large n , the translation length $T_{\tilde{\alpha}_n}$ of $\tilde{\alpha}_n$ satisfies,

$$T_{\tilde{\alpha}_n} \geq \frac{1}{D_n} \log \left(\frac{\varsigma_n e^{D_n T_{\tilde{\alpha}_n^{m_n} \tilde{\beta}_n}} + 3}{\varsigma_n e^{D_n T_{\tilde{\alpha}_n^{m_n} \tilde{\beta}_n}} - 1} \right).$$

This implies that for large n there exists $\epsilon_n \rightarrow 1$ such that,

$$|\tilde{\lambda}_n|^{2D_n} \geq \left(\frac{\epsilon_n \varsigma_n |\text{tr}^{2D_n}(\tilde{\alpha}_n^{m_n} \tilde{\beta}_n)| + 3}{\epsilon_n \varsigma_n |\text{tr}^{2D_n}(\tilde{\alpha}_n^{m_n} \tilde{\beta}_n)| - 1} \right).$$

Hence there exists $M, \rho > 0$ such that for $n > M$, we have $\left| |\tilde{\lambda}_n|^{2D_n} - 1 \right| > \rho |\text{tr}(\tilde{\alpha}_n^{m_n} \tilde{\beta}_n)|^{-2D_n}$, and the lemma follows. $\square$

By Lemma 5.5 and $|\operatorname{tr}(\tilde{\beta}_n)| \asymp |\operatorname{tr}(\beta_n)|(|\lambda_n|^2 - 1)$ we have,

$$\begin{aligned} \lim \frac{1}{|\operatorname{tr}(\tilde{\beta}_n)|(|\tilde{\lambda}_n|^2 - 1)} &\leq \lim \rho(|\operatorname{tr}(\beta_n)|(|\lambda_n|^2 - 1))^{4D_n-1} \\ &\leq \rho'(|\lambda_n|^2 - 1)^{\frac{(1-2D_n)(1-4D_n)}{D_n}} = 0. \end{aligned}$$

Hence we can find Mobius transformations $\varphi_n(x) = \kappa_n x$ such that generators $\varphi_n < \tilde{\alpha}_n, \tilde{\beta}_n > \varphi_n^{-1}$ satisfies conditions of Lemma 4.2.

5.2.2 (A_2)

We can reduce this case to (A_1) , and use similar methods to construct classical generators.

5.3 Case (B)

Set Mobius transformations $\psi_n(x) = \zeta_n \lambda_n^{2k_n-2} x$ and consider the generators $\psi_n < \alpha_n, \alpha_n^{k_n-1} \beta_n > \psi_n^{-1}$ instead we reduce this case to case (A) . Hence we have completed our proof. $\square$

6 Γ with small D_Γ is either classical or existence of universal lower bound on Z_Γ

Lemma 6.1. *There exists $c > 0$ such that: Let Γ_n be a sequence of Schottky groups with $D_n \rightarrow 0$. Then for sufficiently large n , either there exists a subsequence Γ_{n_i} that are classical Schottky groups, or there exists a subsequence Γ_{n_j} with generating set $\langle \alpha_{n_j}, \beta_{n_j} \rangle$ such that $Z_{\langle \alpha_{n_j}, \beta_{n_j} \rangle} > c$.*

Proof. We prove by contradiction. Suppose there exists Γ_n a sequence of Schottky groups such that for every generating set $\langle \alpha_n, \beta_n \rangle$ of Γ_n we have $Z_{\langle \alpha_n, \beta_n \rangle} \rightarrow 0$.

For each n , by replacing $\langle \alpha_n, \beta_n \rangle$ with $\langle \alpha_n, \alpha_n^{m_n} \beta_n \rangle$ for sufficiently large m_n if necessary, we can always assume that every Γ_n is generated by generators with $|\operatorname{tr}(\alpha_n)| \leq |\operatorname{tr}(\beta_n)|$ and $|\operatorname{tr}(\beta_n)| \geq \frac{\log 3}{D_n}$.

We take the upper sapce model $\mathbb{H}^3$. By conjugating with Mobeius transformations, we can assume that α_n have fixed points $0, \infty$ with mutiplier λ_n , and β_n with $z_{\beta_n, u} = 1$. Recall that, as before we denote the two fixed points

of β_n by $z_{\beta_n,l}, z_{\beta_n,u}$, with $|z_{\beta_n,l}| \leq |z_{\beta_n,u}|$. When we write β_n in matrix form, we assume that $|a_n| \leq |d_n|$, otherwise we replace β_n with β_n^{-1} .

By assumption, we have two cases: either case (A) $z_{\beta_n,l} \rightarrow 1$ or case (B) $z_{\beta_n,l} \rightarrow 0$. First we consider the case (A).

6.1 Case (A)

There are two possibilities, (A₁) $\liminf_n |\lambda_n| = 1$ or (A₂) there exists $\lambda > 1$ such that $|\lambda_n| > \lambda$.

6.1.1 (A₂)

Since $|\text{tr}(\beta_n)| \rightarrow \infty$, and $|z_{\beta_n,-} - z_{\beta_n,+}| \rightarrow 0$, there exists Mobius transformations ψ_n such that, for large n , $\psi_n \beta_n \psi_n^{-1}$ have fixed points within $\{z | \frac{1}{\lambda} < |z| < \lambda\}$. Therefore, for sufficiently large n , $\psi_n < \alpha_n, \beta_n > \psi_n^{-1}$ satisfies Lemma 4.2, hence a contradiction.

6.1.2 (A₁)

Take subsequence if necessary, we may assume that $|\lambda_n|$ is strictly decreasing to 1. For large enough n , we choose a sequence of positive integers m_n depends on n such that $1 + \frac{1}{m_n+1} \leq |\lambda_n| \leq 1 + \frac{1}{m_n}$. Let us set $\zeta_n = \frac{a_n}{c_n}, \eta_n = \frac{-d_n}{c_n}$. It follows from Lemma 3.2,

$$|z_{\alpha_n^{m_n} \beta_n, \pm} - \zeta_n \lambda_n^{2m_n}| \leq \rho \frac{|\lambda_n^{m_n}|}{|\text{tr}(\beta_n)|} \leq \rho \frac{e}{|\text{tr}(\beta_n)|}$$

for large n . Also by Lemma 3.2, and the assumption that $z_{\beta_n,l} \rightarrow z_{\beta_n,u} = 1$, we have both $\zeta_n, \eta_n \rightarrow 1$. Hence

$$|z_{\alpha_n^{m_n} \beta_n, \pm} - \lambda_n^{2m_n}| + |z_{\alpha_n^{m_n} \beta_n, \mp} - 1| \rightarrow 0.$$

Since by our choice of m_n , we have $|\lambda_n^{2m_n}| \rightarrow e^2$. It follows that $|z_{\alpha_n^{m_n} \beta_n, \pm}| \rightarrow e^2$, and $|z_{\alpha_n^{m_n} \beta_n, \mp}| \rightarrow 1$. Therefore, there exists $c > 0$ such that $Z_{<\alpha_n, \alpha_n^{m_n} \beta_n>} > c$ for sufficiently large n .

6.2 Case (B)

Here we have either: (B₁) $\liminf_n |\lambda_n| < \Lambda$ for some $\Lambda > 1$, or (B₂) $\liminf_n |\lambda_n| \rightarrow \infty$. We also assume that $|\zeta_n| \leq |\eta_n|$ as before.

6.2.1 (B_1)

Take subsequence if necessary, we may assume that $|\lambda_n| \leq \Lambda$ for large n . Choose positive integers k_n , such that $1 \leq |\zeta_n \lambda_n^{2k_n}| \leq |\lambda_n^2|$. It follows from Lemma 3.2, for $\epsilon > 0$, there is $n > N_\epsilon$ such that $1 - \epsilon < |z_{\alpha_n^{k_n} \beta_n, +}|, |z_{\alpha_n^{k_n} \beta_n, -}| < \Lambda + \epsilon$. There are two possibilities: (B'_1) $\liminf_n |z_{\alpha_n^{k_n} \beta_n, +} - z_{\alpha_n^{k_n} \beta_n, -}| \rightarrow 0$, or (B''_1) $\liminf_n |z_{\alpha_n^{k_n} \beta_n, +} - z_{\alpha_n^{k_n} \beta_n, -}| > 0$. For (B''_1) , we have $Z_{\langle \alpha_n, \alpha_n^{k_n} \beta_n \rangle} > c$, for some $c > 0$ and large n .

Suppose (B'_1) . By passing to subsequence if necessary, we take $|z_{\alpha_n^{k_n} \beta_n, +} - z_{\alpha_n^{k_n} \beta_n, -}| \rightarrow 0$. If $|\lambda_n|$ decreases to 1, then we choose positive integers m_n as defined in case (A_1) . It follows as previous argument, $|z_{\alpha_n^{k_n+m_n} \beta_n, \pm}| \rightarrow e^2$ and $|z_{\alpha_n^{k_n+m_n} \beta_n, \mp}| \rightarrow 1$. Hence there exists $c > 0$, such that $Z_{\langle \alpha_n, \alpha_n^{k_n+m_n} \beta_n \rangle} > c$ for large n .

6.2.2 (B_2)

By taking a subsequence of α_n , we may assume that $|\lambda_n|$ is strictly increasing. Choose a sequence of largest integers $k_n \geq 0$, such that $|\zeta_n \lambda_n^{2k_n}| \leq 1$. If $\limsup |\zeta_n \lambda_n^{2k_n}| = 1$ but $\limsup \zeta_n \lambda_n^{2k_n} \neq 1$, then there is a subsequence $\langle \alpha_{n_j}, \alpha_{n_j}^{k_{n_j}} \beta_{n_j} \rangle$ of $\langle \alpha_n, \alpha_n^{k_n} \beta_n \rangle$, such that $\liminf_j Z_{\langle \alpha_{n_j}, \alpha_{n_j}^{k_{n_j}} \beta_{n_j} \rangle} > 0$.

If $\limsup \zeta_n \lambda_n^{2k_n} = 1$, then let $\langle \alpha_{n_i}, \alpha_{n_i}^{k_{n_i}} \beta_{n_i} \rangle$ be the subsequence of $\langle \alpha_n, \alpha_n^{k_n} \beta_n \rangle$ that is $\lim_i \zeta_{n_i} \lambda_{n_i}^{2k_{n_i}} = 1$.

If $\limsup |\operatorname{tr}(\alpha_{n_i}^{k_{n_i}} \beta_{n_i})| = \infty$, then by pass down to subsequence if necessary, for large i , $\alpha_{n_i}^{k_{n_i}} \beta_{n_i}$ will have disjoint isometric circles. Since $|\operatorname{tr}(\beta_{n_i})| \rightarrow \infty$, and $z_{\beta_{n_i}, u} = 1$ so $|z_{\beta_{n_i}, +} - z_{\beta_{n_i}, -}| \leq 1$, by Lemma 3.2, $\lim_i \min\{|\eta_{n_i} - z_{\beta_{n_i}, -}|, |\eta_{n_i} - z_{\beta_{n_i}, +}|\} \rightarrow 0$. Hence there exists a $\kappa > 0$ such that for large i we have, $\kappa^{-1} < |z_{\alpha_{n_i}^{k_{n_i}} \beta_{n_i}, l}| \leq |z_{\alpha_{n_i}^{k_{n_i}} \beta_{n_i}, u}| < \kappa$. And since $|\lambda_{n_i}| \rightarrow \infty$, we can choose Mobius transformations ψ_i such that $\psi_i \langle \alpha_{n_i}, \alpha_{n_i}^{k_{n_i}} \beta_{n_i} \rangle \psi_i^{-1}$ satisfies Lemma 4.2.

If $\limsup |\operatorname{tr}(\alpha_{n_i}^{k_{n_i}} \beta_{n_i})| < \infty$, then let ϕ_i be Mobius transformations such that $\phi_i \alpha_{n_i}^{k_{n_i}} \beta_{n_i} \phi_i^{-1}$ have fixed points $0, \infty$, and fixed point $z_{\phi_i \alpha_{n_i} \phi_i^{-1}, u}$ of α_{n_i} is 1. Then it follows that $z_{\phi_i \alpha_{n_i} \phi_i^{-1}, l} \rightarrow 1$. Since $|\operatorname{tr}(\phi_i \alpha_{n_i} \phi_i^{-1})| \rightarrow \infty$, hence we have reduced this case to case (A) , which we already considered.

If $\limsup |\zeta_n \lambda_n^{2k_n}| < 1$, then we have two possibilities: (B'_2) $\liminf |\zeta_n \lambda_n^{2k_n+2}| = 1$, or (B''_2) $\liminf |\zeta_n \lambda_n^{2k_n+2}| > 1$.

Consider case (B'_2) . Let $\langle \alpha_{n_i}, \alpha_{n_i}^{k_{n_i}} \beta_{n_i} \rangle$ be a subsequence of $\langle \alpha_n, \alpha_n^{k_n} \beta_n \rangle$

such that $\lim_i |\zeta_{n_i} \lambda_{n_i}^{2k_{n_i}+2}| \rightarrow 1$. If $\sup_i |\operatorname{tr}(\alpha_{n_i}^{k_{n_i}} \beta_{n_i})| < \infty$, then we conjugate $\phi_i \alpha_{n_i} \phi_i^{-1}, \phi_i \alpha_{n_i}^{k_{n_i}} \beta_{n_i} \phi_i^{-1}$ to $\hat{\alpha}_i, \hat{\beta}_i$ respectively with $\hat{\alpha}_i$ have fixed points $0, \infty$ and $\hat{\beta}_i$ have $z_{\hat{\beta}_i, u} = 1$. Since $\sup_i |\hat{\lambda}_i| < \infty$, it follows that $\langle \hat{\alpha}_i, \hat{\beta}_i \rangle$, falls under one of case (A) or case (B_1) . On the other hand, if $\sup_i |\operatorname{tr}(\alpha_{n_i}^{k_{n_i}} \beta_{n_i})| = \infty$, then for large i these will have disjoint isometric circles with radius decreasing to 0. And since the distance between the centers of these isometric circles is decreasing to 0, and also $\inf_i |\lambda_{n_i}|^2 > c > 1$, we can choose Mobius maps $\{\psi_i\}$ that fixes $0, \infty$, so that for large enough i , $\psi_i \alpha_{n_i}^{k_{n_i}} \beta_{n_i} \psi_i^{-1}$ will satisfies Lemma 4.2.

Case (B_2'') . First we define a new sequence of $\langle \tilde{\alpha}_n, \tilde{\beta}_n \rangle$ as follows: Consider $\langle \alpha_n, \alpha_n^{k_n} \beta_n \rangle$. If $|\operatorname{tr}(\alpha_n^{k_n} \beta_n)| \geq |\operatorname{tr}(\alpha_n)|$, then set $\tilde{\alpha}_n = \alpha_n, \tilde{\beta}_n = \beta_n$. Otherwise, let ϕ_n be the Mobius map so that $\phi_n \alpha_n^{k_n} \beta_n \phi_n^{-1}$ have fixed points $0, \infty$, and $\phi_n \alpha_n \phi_n^{-1}$ have $z_{\phi_n \alpha_n \phi_n^{-1}, u} = 1$. Set $\alpha_{n,1} = \phi_n \alpha_n^{k_n} \beta_n \phi_n^{-1}, \beta_{n,1} = \phi_n \alpha_n \phi_n^{-1}$. We define integer $k_{n,1}$ with respect to $\langle \alpha_{n,1}, \beta_{n,1} \rangle$ the same way as we defined k_n before.

Now if $|\operatorname{tr}(\alpha_{n,1}^{k_{n,1}} \beta_{n,1})| \geq |\operatorname{tr}(\alpha_{n,1})|$ then we set $\tilde{\beta}_n = \beta_{n,1}$ and $\tilde{\alpha}_n = \alpha_{n,1}$. Otherwise, we repeat this construction to get a sequence $\langle \alpha_{n,m}, \beta_{n,m} \rangle$. Then there exists a m_n , such that either $|\operatorname{tr}(\alpha_{n,m_n}^{k_{n,m_n}} \beta_{n,m_n})| > |\operatorname{tr}(\alpha_{n,m_n})| - \frac{1}{n}$, or $|\operatorname{tr}(\alpha_{n,m_n})| < \frac{1}{n}$. We define $\tilde{\alpha}_n = \alpha_{n,m_n}, \tilde{\beta}_n = \beta_{n,m_n}$.

Now consider $\langle \tilde{\alpha}_n, \tilde{\beta}_n \rangle$. If $\liminf_n |\operatorname{tr}(\tilde{\alpha}_n)| < \infty$, we choose a subsequence with $|\operatorname{tr}(\tilde{\alpha}_{n_i})| < c$ for all large i and some $c > 0$. Let p_i be a sequence of least positive integers such that $|\operatorname{tr}(\tilde{\alpha}_{n_i}^{p_i} \tilde{\beta}_{n_i})| > \frac{1}{D_{n_i}}$. We conjugate $\tilde{\alpha}_{n_i}^{p_i} \tilde{\beta}_{n_i}$ by ψ_i that fixes $0, \infty$ and $z_{\psi_i \tilde{\alpha}_{n_i}^{p_i} \tilde{\beta}_{n_i} \psi_i^{-1}, u} = 1$. Set $\bar{\alpha}_i = \psi_i \tilde{\alpha}_{n_i} \psi_i^{-1}, \bar{\beta}_i = \psi_i \tilde{\alpha}_{n_i}^{p_i} \tilde{\beta}_{n_i} \psi_i^{-1}$. By construction, $\langle \bar{\alpha}_i, \bar{\beta}_i \rangle$ satisfies either case (A) or case (B_1) . In either case, we are done.

On the other hand, suppose $\liminf_n |\operatorname{tr}(\tilde{\alpha}_n)| = \infty$. If $\liminf_n |\operatorname{tr}(\tilde{\beta}_n)| < \infty$, then let ϕ_n be Mobius maps such that $\phi_n \tilde{\beta}_n \phi_n^{-1}$ have fixed points $0, \infty$ and $z_{\phi_n \tilde{\alpha}_n \phi_n^{-1}, u} = 1$. Then $\langle \phi_n \tilde{\beta}_n \phi_n^{-1}, \phi_n \tilde{\alpha}_n \phi_n^{-1} \rangle$ satisfies either case (A) or (B_1) , hence we are done. If $\liminf_n |\operatorname{tr}(\tilde{\beta}_n)| = \infty$, its sufficient to assume that $\langle \tilde{\alpha}_n, \tilde{\beta}_n \rangle$ satisfies case (B_2'') , otherwise we are done. We define $\tilde{k}_n = k_{n,m_n}$.

First we claim that

$$\liminf_n \frac{|\zeta_n \tilde{\lambda}_n^{2\tilde{k}_n}| - |\tilde{\lambda}_n|^{-2}}{|\tilde{\lambda}_n|^{-1}} = \infty.$$

Otherwise, there exists $\sigma > 0$, and a subsequence such that

$$\lim_i \frac{|\tilde{\zeta}_{n_i} \tilde{\lambda}_{n_i}^{2\tilde{k}_{n_i}}| - |\tilde{\lambda}_{n_i}^{-2}|}{|\tilde{\lambda}_{n_i}^{-1}|} < \sigma.$$

Set $\nu_i = \tilde{\alpha}_{n_i}^{\tilde{k}_{n_i}} \tilde{\beta}_{n_i}$, $\mu_i = \tilde{\alpha}_{n_i}$. Since $|\operatorname{tr}(\tilde{\beta}_n)| \geq |\operatorname{tr}(\tilde{\alpha}_n)|$ and $|z_{\tilde{\beta}_n, -} - z_{\tilde{\beta}_n, +}| \leq 1$, it follows from Lemma 3.2 and $\lim_i \frac{|\operatorname{tr}(\nu_i)|}{|\tilde{\lambda}_{n_i}|} > \epsilon$ for $\epsilon > 0$, we have $|\eta_{\nu_i} - 1| < \delta |\tilde{\lambda}_{n_i}|^{-2}$, for some $\delta > 0$. Hence $|\eta_{\nu_i \mu_i} - \tilde{\lambda}_{n_i}^{-2}| < \delta |\tilde{\lambda}_{n_i}^{-4}|$, for large i . Let ϕ_i be Mobius transformations define by $\phi_i(x) = \sqrt{\tilde{\zeta}_{n_i} \tilde{\lambda}_{n_i}^{2\tilde{k}_{n_i}}} x$ for $x \in \mathbb{C}$. Set $\nu_{i,\phi} = \phi_i \nu_i \phi_i^{-1}$, $\mu_{i,\phi} = \phi_i \mu_i \phi_i^{-1}$. Then for large i ,

$$|\eta_{\nu_{i,\phi} \mu_{i,\phi}}| > \frac{1}{\sigma |\tilde{\lambda}_{n_i}|^{-1} + 1} - \frac{\delta}{\sigma |\tilde{\lambda}_{n_i}|^3 + |\tilde{\lambda}_{n_i}|^2}.$$

Since $|\tilde{\zeta}_{n_i} \tilde{\lambda}_{n_i}^{2\tilde{k}_{n_i}}| > |\tilde{\lambda}_{n_i}|^{-2}$, we also have

$$|\eta_{\nu_{i,\phi} \mu_{i,\phi}}| < 1 + \frac{\delta}{|\tilde{\lambda}_{n_i}|^2}.$$

Hence, $|\eta_{\nu_{i,\phi} \mu_{i,\phi}} - \zeta_{\nu_{i,\phi} \mu_{i,\phi}}| < 1 + \epsilon |\tilde{\lambda}_{n_i}|^{-1}$, for large i and $\epsilon > 0$. Note that, $\zeta_{\nu_{i,\phi} \mu_{i,\phi}} = 1$. If $\liminf_i |\operatorname{tr}(\nu_{i,\phi} \mu_{i,\phi})| = \infty$, then the radius of isometric circles $\Re_{\nu_{i,\phi} \mu_{i,\phi}}$ of $\nu_{i,\phi} \mu_{i,\phi}$ well decrease to 0. Since $\lim_i |\tilde{\lambda}_{n_i}| = \infty$, we can choose Mobius transformations ψ_i such that $\psi_i \nu_{i,\phi} \mu_{i,\phi} \psi_i^{-1}$ will have fixed points within the annulus $\{z \in \mathbb{C} | \kappa^{-1} < |z| < \kappa\}$, for some $\kappa > 0$. Hence $\psi_i < \mu_{i,\phi}$, $\nu_{i,\phi} \mu_{i,\phi} > \psi_i^{-1}$ satisfies Lemma 4.2 for large i .

On the other hand, If $\liminf_i |\operatorname{tr}(\nu_{i,\phi} \mu_{i,\phi})| < \infty$, then choose Mobius transformations ψ_i such that, $\psi_i \nu_{i,\phi} \mu_{i,\phi} \psi_i^{-1}$ have fixed points 0, ∞ and $z_{\mu_{i,\phi}, u} = 1$. Then $\phi_i < \nu_{i,\phi} \mu_{i,\phi}$, $\mu_{i,\phi} > \phi_i^{-1}$ satisfies either case (A) or case (B_1), hence we are done. This completes our proofs of the claim.

By our claim, there exists $0 < \rho_n \rightarrow \infty$, such that $|\tilde{\zeta}_n \tilde{\lambda}_n^{2\tilde{k}_n}| - |\tilde{\lambda}_n|^{-2} > \rho_n |\tilde{\lambda}_n|^{-1}$. From the construction of $\tilde{\alpha}_n$ and $\tilde{\beta}_n$, it follows that $|\operatorname{tr}(\tilde{\alpha}_n^{\tilde{k}_n} \tilde{\beta}_n)| > c |\operatorname{tr}(\tilde{\alpha}_n)|$, for $c > 0$ and all large n . Hence there exists $\kappa > 0$, $|\operatorname{tr}(\tilde{\alpha}_n^{\tilde{k}_n} \tilde{\beta}_n)|^{-1} < \kappa |\tilde{\lambda}_n|^{-1}$. Therefore, $|\tilde{\zeta}_n \tilde{\lambda}_n^{2\tilde{k}_n}| - |\tilde{\lambda}_n|^{-2} > \rho_n \kappa^{-1} |\operatorname{tr}(\tilde{\alpha}_n^{\tilde{k}_n} \tilde{\beta}_n)|^{-1}$. Let χ_n be Mobius transformations defined by $\chi_n(z) = |\tilde{\lambda}_n| z$. Then it follows from these estimates that for large n , $\chi_n < \tilde{\alpha}_n$, $\tilde{\alpha}_n^{\tilde{k}_n} \tilde{\beta}_n > \chi_n^{-1}$ have fixed points within the annulus $\{z \in \mathbb{C} | |\tilde{\lambda}_n|^{-1} < |z| < |\tilde{\lambda}_n|\}$ and satisfies Lemma 4.2, and this complete our proof. $\square$

7 Proof of Main Theorem

Theorem 7.1. *There exists $\epsilon > 0$ such that every 2-generated Schottky group Γ with $D_\Gamma < \epsilon$ is a classical Schottky group.*

Proof. This follows from Theorem 5.1 and Lemma 6.1. $\square$

Proof of Theorem 1.1. Let Γ' be a nonelementary finitely generated Kleinian group. Selberg lemma implies Γ' contains a torsion-free subgroup Γ'' of finite index, in particular $\mathfrak{D}_{\Gamma'} = \mathfrak{D}_{\Gamma''}$. Note that if Γ'' is geometrically infinite with $\Omega_{\Gamma''} \neq \emptyset$ then $D_{\Gamma''} = 2$, this implies $\mathfrak{D}_{\Gamma''} = 2$ for geometrically infinite groups. So we can assume Γ'' is geometrically finite. If Γ'' contains parabolic of rank $l_{\Gamma''}$ then $D_{\Gamma''} \geq l_{\Gamma''}/2$. Hence, for sufficiently small Hausdorff dimension $D_{\Gamma''}$, we can assume Γ'' is convex-cocompact of second kind.

It follows from Ahlfors finiteness theorem, that $\Omega_{\Gamma''}/\Gamma''$ consists of finite number of compact Riemann surfaces. Let S be a component of $\Omega_{\Gamma''}/\Gamma''$. If S is incompressible then $\pi_1(S)$ is a surface subgroup of Γ'' . Since $1 = D_{\pi_1(S)} \leq D_{\Gamma''}$, if $D_{\Gamma''}$ is small, we may assume S is compressible. Then we can decompose Γ'' along the compression disk. After repeating the decomposition process finitely many times we are left with topological balls, i.e. $\mathbb{H}^3/\Gamma''$ is a handle body. This implies Γ'' is a finitely generated free purely loxodromic Kleinian group of second kind, i.e. Γ'' is a Schottky group.

By assuming the limit set have sufficiently small Hausdorff dimension we have reduced the general case to the case of Schottky groups. Now it follows from Marden's rigidity theorem, all Schottky groups of the same rank are quasiconformally equivalent. Therefore we have from Theorem 7.1, there exists $\lambda > 0$ such that all nonelementary finitely generated Kleinian Γ' with $\mathfrak{D}_{\Gamma'} \leq \lambda$ contains a classical Schottky group of finite index. It follows that we have a strict lower bound on the Hausdorff dimension of all non-classical Schottky group. $\square$

ACKNOWLEDGEMENTS

I am indebted to Ying Zhou for all the support and unwavering help that has been provided to me during this work. I would like to express my gratitude to Peter Shalen, Dick Canary, and Marc Culler for their time spend, interests, and valuable suggestions on the paper.

E-mail: yonghou@math.uic.edu

References

- [1] Anderson, J., Canary, R., Culler, M., Shalen, B.P. (1996). Free Kleinian groups and volumes of hyperbolic 3-manifolds. *J. Diff. Geo.*, 44, 738-782.
- [2] Ancona, A. (1990). Théorie du potentiel sur les graphes et les variétés. *Lect. Notes in Math.* 1427, Springer, Berlin.
- [3] A. Berdon. (1983). *The Geometry of discrete groups*, Springer-verlag.
- [4] Bishop, C. Jones, P. (1997). Hausdorff dimension and Kleinian groups. *Acta. Math.* 179. No 1, 1-39.
- [5] Canary, D. Taylor, E. (1994). Kleinian groups with small limit sets. *Duke Mathematical Journal.* 73, 371-381.
- [6] Chuckrow, V. (1968). On Schottky groups with applications to Kleinian groups. *Ann. Math.* 88. 47-61
- [7] Culler, M., Shalen, B.P. (1992). Paradoxical decompositions, 2-generator kleinian groups, and volumes of hyperbolic 3-manifolds. *J. Amer. Math. Soc.* 5, 231-288.
- [8] Culler, M., Hersonsky, S., Shalen, B.P. (1998). The first Betti number of the smallest closed hyperbolic 3-manifolds. *Topology.* Vol. 37 No 4. pp 805-849.
- [9] Doyle, P. (1988) On the bass note of a Schottky group. *Acta Math.* 160, 249-284.
- [10] Gehring, M., Maclachlan, M. (1998). Two-generator arithmetic Kleinian groups II, *Bull. Lond. Math. Soc.*, 30, N 3, 258-266.
- [11] Gromov, M. (1987). Hyperbolic Groups. In *essays in group theory*. Gersten (editor) M.S.R.I. Publ. 8. Springer, Berlin. pp 75-263.
- [12] Hou, Y. (2001). Critical exponent and displacement of negatively curved free groups. *J. Diff. Geo.* 57, 173-195.
- [13] Kapovich, M. (2000). *Hyperbolic Manifolds and Discrete Groups: Lecture on Thurston's Hyperbolization*, Birkhauser's series "Progress in mathematics".

- [14] Marden, A. (1974). Schottky groups and circles. Contributions to analysis (a collection of papers dedicated to Lipman Bers). pp. 273-278. Academic Press.
- [15] Patterson, S.J. (1987). Measures on limit sets of Kleinian groups. Analytical and Geometrical Aspects of hyperbolic space. Cambridge Univ. Press. pp 291-323.
- [16] Phillips, R., Sarnak, P. The Laplacian for domains in hyperbolic space and limit sets of Kleinian groups. Acta Math., 155:173-241
- [17] Sullivan, D. (1982). Discrete conformal groups and measurable dynamics. Bull. Amer. Math. Soc. 6, 57-73.